

## Format of ACE solar wind data plots (produced by Ian Richardson, GSFC, [richardson@lheavx.gsfc.nasa.gov](mailto:richardson@lheavx.gsfc.nasa.gov))

The plots show 64s resolution “level 2” (verified) magnetic field and plasma data from the ACE MAG and SWEPAM experiments, except for events in 1997 and from July 2004, when 5-minute “browse” (preliminary) data are shown. The data are obtained from the ACE Science Center (<http://www.srl.caltech.edu/ACE/ASC/>). For each event, a 2-day interval is shown, centered on the time of peak Dst as noted in the CDAW event table. The parameters shown are:

- The magnetic field intensity, polar (90 deg = northward) and azimuthal (90 deg = directed to the west) angles (GSE);
- Proton temperature (black line). The red line is the “expected temperature” for normally expanding solar wind (Richardson and Cane, 1995). This is derived from the solar wind speed (panel 6) using the results of Lopez (1987) based on OMNI data. See Neugebauer et al. (2003) for a similar formula derived from ACE data. In normal (non-ICME) solar wind,  $T_{\text{exp}}$  typically tracks  $T_p$ . In ICMEs, typically  $T_p \ll T_{\text{exp}}$ . Black shaded regions indicate when  $T_p < 0.5T_{\text{exp}}$ , a frequent indicator of possible ICMEs (Richardson and Cane, 1995), as well as other structures such as heliospheric current/plasma sheet crossings.
- Solar wind proton density;
- Solar wind speed;
- Solar wind helium/proton ratio. Elevated values above 0.06 are typically associated with ICMEs. However, not all ICMEs have elevated He/p. This ratio also varies during the solar cycle (see Richardson and Cane, 2004, and references therein);
- Ratio of solar wind oxygen ions with charges 7 and 6, from the ACE/SWICS instrument (Level 2 data from the ACE Science Center).  $O^7/O^6$  is believed to provide a measure of the coronal electron freezing-in-temperature. The red line is an estimate of the expected value of  $O^7/O^6$  for ambient, non-ICME solar wind, inferred from the solar wind speed (Richardson and Cane, 2004, and references therein). ICMEs typically have  $O^7/O^6 \gg$  the expected value;
- (1998 – 2000 events) Fraction of solar wind iron ions with charge states  $\geq 16$ , from ACE/SWICS. High values ( $>0.1$ ) are typically associated with ICMEs (e.g., Lepri et al., 2001). Red line indicates “expected value” based on the solar wind speed (Richardson & Cane, 2004). (Data supplied by Sue Lepri, U. of Michigan);
- Bz in GSM coordinates. Negative values indicate southward-directed magnetic fields;
- The Dst geomagnetic index;
- The Kp geomagnetic index, shown as Kp\*10;
- An ICME index (1 = probable ICME), based on the ICME identifications of Cane and Richardson (2003), with updates. Note that these identifications did not use composition/charge state data, thus some adjustments to the event boundaries may be made in the light of these data. In addition, it is possible that an occasional event may have been missed or mis-identified.

Vertical green lines indicate times of *geomagnetic storm sudden commencements* with at least one “A” report in the NGDC list. These typically are associated with shocks

observed at ACE, with a delay due to the upstream location of ACE at the L1 libration point. Other shocks may be evident in the plasma/field data - see the ACE shock list at [http://www-ssg.sr.unh.edu/mag/ace/ACElists/obs\\_list.html#shocks](http://www-ssg.sr.unh.edu/mag/ace/ACElists/obs_list.html#shocks).

#### References:

H. V. Cane and I. G. Richardson, "Interplanetary coronal mass ejections in the near-Earth solar wind during 1996-2002", *J. Geophys. Res.*, 108 No. 4, 10.1029/2002JA009817, 2003.

S. T. Lepri, T. H. Zurbuchen, L. A. Fisk, I. G. Richardson, H. V. Cane, and G. Gloeckler, "Iron charge distribution as an identifier of interplanetary coronal mass ejections", *J. Geophys. Res.*, 106, 29,231, 2001.

Lopez, R. E., Solar cycle invariance in solar wind proton temperature relationships, *J. Geophys. Res.*, 92, 11,189, 1987.

M. Neugebauer et al., Genesis on-board determination of the solar wind flow regime, *Space Sci. Rev.* 105, 661, 2003.

I.G. Richardson and H.V. Cane, "Regions of abnormally low proton temperature in the solar wind (1965-1991) and their association with ejecta", *J. Geophys. Res.*, 100, 23397, 1995.

I. G. Richardson and H. V. Cane, "Identification of interplanetary coronal mass ejections at 1 AU using multiple solar wind composition anomalies", *J. Geophys. Res.*, 109, A09104, doi:10.1029/2004JA010598, 2004.